



The imprint of windblown dust from the North American Southwest on the California Channel Islands and Pacific Ocean sediments

G.E. Jardine^a, A.J. Crocker^{a, *}, I. Bailey^b, M.J. Cooper^a, J.A. Milton^a, P.A. Wilson^{a, **}

^a University of Southampton, Waterfront Campus, National Oceanography Centre, Southampton, SO14 3ZH, UK

^b Camborne School of Mines and Environment and Sustainability Institute, University of Exeter, Penryn Campus, Cornwall, UK

ARTICLE INFO

Article history:

Received 19 October 2020

Received in revised form

9 March 2021

Accepted 4 April 2021

Available online xxx

Handling Editor: I Hendry

Keywords:

Present

Quaternary

Paleoclimatology

North America

Radiogenic isotopes

ABSTRACT

Climate projections for the North American Southwest (NASW) predict an increasing frequency and duration of droughts over the 21st century in response to human-induced warming, with potentially severe economic and social consequences. The geological record provides a way to contextualise this prediction because of the past occurrence of abrupt hemispheric warming events and sustained intervals of atmospheric carbon dioxide loading equivalent to those projected for AD 2100 (between ~500 and 900 ppmv). Yet, terrestrial climate archives are typically too short and incomplete to provide a full record of these events. In principle, drill cores from deep sea sediments in the eastern Pacific Ocean can be used to overcome this problem because they contain long records of continental dust and distal riverine-supplied sediments from North America. Yet our limited understanding of the provenance and transport pathways of these sediments impedes use of these marine archives for this purpose. Here we present radiogenic isotope data (Sr, Nd and Pb) from known NASW dust-producing hot spots – playa lakes in the Mojave Desert, Quaternary silts mantling the California Channel Islands and the terrigenous fraction from marine sediments of the eastern Pacific Ocean, supported by new maps of bedrock isotopic composition in the NASW. We use these and published data sets to infer the origin of playa lake silts in the Mojave Desert and the source of windblown sediments to the California Channel Islands and nearby ocean basins. Our Results rule out a significant contribution from the distal tails of either the Pacific Asian dust plume or the North African dust plume to the Quaternary Channel Island silt mantles, corroborating the suggestion that they are aeolian in origin and sourced from the NASW on the Santa Ana winds. We identify the Outer California Borderland basins as an attractive proposition for studying past dust flux and palaeoaridity in the North American Southwest.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Climate projections to the year 2100 predict an increased frequency and duration of droughts in response to human-induced warming in mid-latitude arid to semi-arid regions (Ault et al., 2016; Balling and Goodrich, 2010; Cayan et al., 2010; Cook et al., 2015; Seager et al., 2007). In the North American Southwest (NASW; typically taken to include the states of California, Nevada, Utah, Colorado, New Mexico and Arizona, Fig. 1a), these changes are already evident (Ault et al., 2016; Balling and Goodrich, 2010; Cook et al., 2015; Seager et al., 2007). The 2012–2016 Californian drought

cost the agriculture industry an estimated \$603 million and the loss of over 1500 jobs in 2016 alone (Medellín-Azuara et al., 2016). There is a pressing need, therefore, to assess the robustness of model predictions for hydroclimate in this region (Cook et al., 2015; MacDonald et al., 2008).

Past climate records help to ground-truth future climate projections and geological data provide a way to extend beyond the modern observational record by reaching deeper into the past to cover a greater range of past climate conditions. Dendrochronological reconstructions document numerous multi-decadal ‘mega’ droughts during the past 2000 years (Cook et al., 2007, 2014; Salzer and Kipfmüller, 2005). Although long lasting and severe (decadal in length, covering at least 50% of the North American west), these megadroughts occurred when atmospheric carbon dioxide levels were substantially lower than today (~280 ppmv during the Medieval Warm Period, compared with ~415 ppmv, January 2021 (Ahn

* Corresponding author.

** Corresponding author.

E-mail addresses: anya.crocker@noc.soton.ac.uk (A.J. Crocker), paw1@noc.soton.ac.uk (P.A. Wilson).




Fig. 1. Study region with locations of new and previously published data reported herein: a) North American states typically included in the North American Southwest (NASW) (clockwise from left: California, Nevada, Utah, Colorado, New Mexico, Arizona) with five major deserts highlighted (Google Earth imagery). Locations of DSDP sites 469 and 32, ODP sites 893 and 1015 and the sampled Mojave Desert playa lakes (yellow filled circles) also shown. Blue box in the lower left-hand corner denotes area shown in the next panel. b) MODIS Satellite imagery showing dust plumes transported over the Californian Borderland Basins by the Santa Ana wind system on 9th February 2002 (NASA, 2002). c) Map of southern California topography and offshore bathymetry showing location of Inner and Outer Borderland Basins, the coastal Transverse and Peninsular mountain ranges, the locations of ODP sites 893 and 1015 and locations of Anacapa and San Clemente Islands. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

et al., 2012; Keeling et al., 2001)) and only about one third as high as the levels predicted for 2100 under a 'business as usual' emissions scenario (>900 ppm under RCP 8.5 (Meinshausen et al., 2011)). It is instructive, therefore, to reach deeper into the geological past, to study intervals that incorporate the range of estimates for 21st century greenhouse gas levels such as the mid-Piacenzian Warm Period (3.3–3 Ma) (e.g. de la Vega et al., 2020) and Miocene (e.g.

Super et al., 2018).

Lake sediments provide valuable archives of terrestrial hydro-climate in the NASW, including records of variability in aeolian dust and plant leaf waxes (Feakins et al., 2014; Routson et al., 2016; Tchakerian and Lancaster, 2002), but in dryland and arid-prone regions, lakes are short-lived and age control can be challenging. To study pre-Quaternary palaeoclimate states, we turn to marine

sediment core archives, which have the potential to provide continuous, long, well-dated records of terrestrial hydroclimate in deeper time (e.g. Rea, 1994; Tiedemann et al., 1994). Mineral dust deposition in sediment cores can be used as a proxy for aridity in the terrestrial source region because of the strong (but not necessarily simple) relationship between precipitation and natural dust emissions (Hoell et al., 2014; Just et al., 2012; Larrasoana et al., 2003; Nizou et al., 2011; Pettke et al., 2000; Pye, 1987b; Seo et al., 2014, 2015). Yet, marine sediment cores may also receive terrestrial sediments sourced from rivers or ocean currents. Terrigenous flux to the marine realm cannot, therefore, always simply be interpreted as an indicator of aridity, even when those terrigenous sediments are sourced from regions that are dry today. To decipher better the climate record imprinted on sediment cores we need to distinguish between different terrigenous sources and different modes of delivery to the ocean.

Radiogenic isotopes, particularly lead (Pb), neodymium (Nd) and strontium (Sr) are valuable tools to help distinguish between the provenance of sedimentary material accumulating in marine settings over a wide range of spatial and temporal scales (e.g. Abouchami and Zabel, 2003; Cole et al., 2009; Hyeong et al., 2011; Nakai et al., 1993; Pettke et al., 2000; Seo et al., 2014; Xie and Marcantonio, 2012). One particularly influential study of core-top sediments characterised pelagic clays delivered to the central and eastern Pacific Ocean as a mixture of three primary detrital components derived from continental source areas in Asia, Central and South America and North America (Stancin et al., 2006). The fidelity of these interpretations is weakened, however, by limited information on NASW sources.

Here we present (i) new radiogenic isotope data (Pb, Nd and Sr isotopic compositions) for dust source regions in the NASW, silt mantles from the Channel Islands offshore southern California and the detrital silicate fraction of young marine sediment samples from Ocean Drilling Program (ODP) sites 1015 and 893 (Fig. 1a,c) and (ii) new maps of bedrock isotopic composition in the NASW. We use these new data, together with published data, to improve understanding of the geochemical fingerprints of terrigenous sediments transported to the Pacific Ocean from the North American continent.

2. Materials and methods

2.1. Sample selection and description

To investigate the export of terrigenous sediments from North America to the Pacific Ocean from source to sink, we studied three main types of samples: silt mantles from two California Channel Islands (Anacapa and San Clemente) proposed to derive from airborne dust from the North American mainland (Muhs et al., 2007, 2008), surface sediments from playa lakes located in the Mojave Desert and marine sediments from two ODP sites (1015 and 893) on the California margin (Fig. 1a,c). ODP Site 1015 is located in the Santa Monica Basin (33°42.925'N, 118°49.185'W; Fig. 1c) at 901 m below sea level (mbsl); ODP Site 893 is located in the Santa Barbara Basin (34°17.25'N, 120°02.19'W; Fig. 1c) at 576.5 mbsl. The Mojave Desert and island samples were obtained from the United States Geological Survey (USGS), Colorado (courtesy of D. Muhs).

The Channel Island silt mantles are described by Muhs et al. (2007, 2008) as massive, silt-rich horizons that resemble loess, or aeolian silt, ranging from 2 to 30 cm thick, which contrast sharply with the lower subsoil. Precise dating of the silt mantles on San Clemente is difficult but they drape the youngest marine terrace which gives a maximum age of ~80 ka, although they could possibly be younger than 3 ka, based upon a single radiocarbon data point (Muhs et al., 2007). Mineralogical, geochemical and grain size

profile measurements (Muhs et al., 2007, 2008) suggest that the dominant source of the island silt mantles is from the North American continent (not from the island bedrock), implying a windblown origin, but the influence of an Asian dust source (Muhs et al., 2007) and perhaps the Sahara cannot be ruled out. Muhs et al. (2007) and Muhs et al. (2008) also report that the geochemical fingerprint of the fine fraction (<2 µm; clay) in these silt mantles differs slightly from that of the coarser (2–53 µm; silt sized) fraction of the samples, hinting that the silts and clays within the silt mantles may have different provenances. The identity of the North American source is equivocal – both the Mojave Desert and the coastal Transverse and Peninsular mountains of Southern California have been considered as potential source regions for this wind-blown material (Muhs et al., 2007, 2008).

To better characterise sources of aeolian supply to the Californian borderland from the interior of the North American continent we targeted silts from the playa lakes of the Mojave Desert. Playa lake and river valley sediments are known to be active sources of silt-sized material for deflation and aeolian transport in dry conditions (Pye, 1987b) and offshore dust transport on the Santa Ana winds that blow from the inland deserts through the mountain passes of Southern California in response to high pressure systems over the Great Basin can be observed in satellite imagery (Muhs et al., 2007, Fig. 1b).

In addition to wind-borne transport of lithogenic material, the Santa Barbara and Santa Monica basins also receive fluvial sediment eroded and transported by the Santa Clara and Ventura River systems from the Transverse Ranges (Fig. 1c), with the Santa Clara River system supplying more than three times the amount of riverine material to the Santa Barbara and inner borderland basins as the next largest source (Inman and Jenkins, 1999). We sampled sediments of Holocene and deglacial age from ODP sites 1015 and 893 (locations Fig. 1c) to fingerprint these fluvial inputs. “Riverine” samples were selected to characterise the isotopic composition of fluvially derived sediments delivered to the offshore borderland basins from the Californian coastal river valleys during the Late Quaternary. These consist of macroscopically obvious turbidites ultimately originating from fluvial sources at Site 1015 and conspicuous cm-scale ‘grey-layers’ inferred to represent fine-grained flood deposits at Site 893 (Behl, 1995; Marsaglia et al., 1995). “Hemipelagic” samples were also selected at ODP Site 1015, targeting intervals of background sedimentation in between the event horizons.

2.2. Isotopic analyses

The Channel Island silt mantle sediments were sieved to remove the >63 µm fraction, following Muhs et al. (2007, 2008). A subset of these samples was also sieved over a fine mesh to isolate the <10 µm fraction to allow comparison with long-range transported Asian material. Prior to full chemical digestion, carbonates, organic material, biogenic silica and Fe–Mn oxides were removed from the fine fraction of the sieved Channel Island samples and bulk Mojave Desert playa lake silts and Californian borderland ODP sediment samples by a series of chemical preparation steps, adapted from Bayon et al. (2002). Approximately 5 g of dry bulk sediment was placed in sealable Nalgene conical flasks. A solution of ~10–25% acetic acid solution was used to decarbonate the samples, and then 10% hydrogen peroxide solution at 60 °C was used to remove organic matter. Biogenic silica was removed from the Channel Island silt mantles and Californian borderland ODP sediments using a 1.5M NaOH solution following the method of Povea et al. (2015). Fe and Mn oxides were removed from the silt mantle and borderland basin sediments using a 0.05 M hydroxylamine hydrochloride – 15% acetic acid – 0.03 M Na-EDTA solution buffered to pH4 with

analytical grade NaOH in two steps: an initial 3 h reaction period; followed by the replacement of the reagent with fresh solution and a second reaction period of 24 h. For both these steps, the samples were left on a shaker table. For the Mojave Desert playa lake samples, the Fe–Mn removal step was not deemed necessary because tests showed no significant isotopic effect from an anthropogenic contribution of Pb from a subset of samples processed with a Fe–Mn oxide removal step.

For Pb, Nd and Sr isotopic analyses, the chemically processed sediment samples were freeze-dried and approximately 50 mg of freeze-dried sample was weighed into 15 ml Teflon pots. The samples were treated with a final additional cleaning step of 1M ammonium acetate solution following [Pettke et al. \(2000\)](#). Concentrated hydrofluoric acid (HF, ~27M) and concentrated nitric acid (HNO₃, ~12M) were added in a 5:1 ratio and the pots left to reflux at ~130 °C for 48 h. Pb and Nd were then separated from the digested mother solution by ion exchange column procedures and ion selective resin for Sr. Where necessary, a modified column sequence with an additional ion exchange column was used to enhance the separation of Ba from the Sr fraction. Average digestion blanks for Pb, Nd and Sr were 320 pg, 160 pg and 7 ng respectively. One digestion blank (affecting five samples) contained an anomalous concentration of Pb and was excluded from this calculation but review of the data from those five samples indicates no obvious offset from the remainder of the dataset (see [Supplementary Fig. 1](#)), so they are included herein. Average column blanks for Pb, Nd and Sr were 168 pg, 2 pg and 13.5 pg respectively. The Nd-isotope (¹⁴³Nd/¹⁴⁴Nd) and Pb-isotope ratios (²⁰⁶Pb/²⁰⁴Pb, ²⁰⁷Pb/²⁰⁴Pb and ²⁰⁸Pb/²⁰⁴Pb) of the samples were measured at the University of Southampton using a multi-collector inductively coupled plasma mass spectrometer (MC-ICP-MS, Thermo Scientific Neptune). Pb isotopic compositions were analysed using the double spike method of [Taylor et al. \(2015\)](#). The average values of NIST 981 were 16.9394 ± 0.0025 , 15.4955 ± 0.0026 and 36.7093 ± 0.0078 for ²⁰⁶Pb/²⁰⁴Pb, ²⁰⁷Pb/²⁰⁴Pb and ²⁰⁸Pb/²⁰⁴Pb respectively. Nd isotopic compositions were adjusted to a ¹⁴⁶Nd/¹⁴⁴Nd value of 0.7219 ([Vance and Thirlwall, 2002](#)). Mass-bias corrected ratios were normalized to the given ¹⁴³Nd/¹⁴⁴Nd value (0.512115) of the standard JNdi-1 ([Tanaka et al., 2000](#)). The average value of JNdi-1 run as unknowns were 0.512114 ± 0.000006 . ¹⁴³Nd/¹⁴⁴Nd ratios are reported in epsilon notation:

$$\epsilon_{Nd} = \left(\frac{{}^{143}\text{Nd}/{}^{144}\text{Nd}_{\text{sample}}}{{}^{143}\text{Nd}/{}^{144}\text{Nd}_{\text{CHUR}}} - 1 \right) * 10^4 \quad (1)$$

where ¹⁴³Nd/¹⁴⁴Nd_{CHUR} is the Chondritic Uniform Reservoir value of 0.512638 ([Jacobsen and Wasserburg, 1980](#)).

The ⁸⁷Sr/⁸⁶Sr isotopic composition of the samples was measured at the University of Southampton using a thermal ionization mass spectrometer (Thermo-Fisher TRITON Plus) with an ⁸⁸Sr beam of 2V. Fractionation was corrected using an exponential correction normalised to ⁸⁶Sr/⁸⁸Sr = 0.1194. NIST 987 was run as a reference standard, with a long-term mean value of 0.710243 ± 0.000021 (2σ) reported from repeat analyses on this instrument.

2.3. Radiogenic isotope base maps

To place our data and the published radiogenic isotope datasets of NASW dust sources into a regional context we developed Pb, Nd and Sr isotope maps of the bedrock geology by compiling published measurements from the scientific literature (full reference list is given in [Supplementary Information](#)). Most of the data used to develop these maps are for whole rock samples, but for some data accessed through the North American Volcanic and Intrusive Rock

Database (NAVDAT, navdat.org), this information was not available.

All data were assigned latitude and longitude coordinates using published site location information in conjunction with Google Earth (format X°X'X"). The location information was then converted to decimal degrees and input into MATLAB_R2017a. A subset of the latitude and longitude coordinates was derived by generating a path function within Google Earth and using the MATLAB script `kml_shapefile.m` ([Toomey, 2010](#)) to convert the resulting.kml file to a MATLAB shape file that contained the coordinate information.

To generate the colour-shaded map background, the geo-referenced data were averaged into 0.125° × 0.125° latitude/longitude grid squares within the bounds of 110 to 125°W and 30 to 40°N. This irregularly spaced data set of grid-square-averaged isotope data points was used to generate a regularly spaced 3D gridded data set, with the x and y coordinates given by the latitude and longitude, and the z coordinate the associated isotopic value using the function `griddata.m`. This function generated an isotopic 'data' value for the empty spaces in the original data distribution using a natural interpolation function. Some samples of granitoid rocks have much more radiogenic Sr values than typical rocks in the NASW, resulting in a skewed distribution of ⁸⁷Sr/⁸⁶Sr bedrock data. As such, the grid-averaged dataset was filtered to exclude ⁸⁷Sr/⁸⁶Sr values > 0.7350 to distinguish regional trends free of extreme local effects. ²⁰⁶Pb/²⁰⁴Pb isotope values > 21 were also excluded for similar reasons. Original, unfiltered maps are shown in [Supplementary Fig. 2](#). The interpolation function does not allow for an extrapolation outside the area bounded by data points, resulting in some regions in the map that lack a colour-coded background. The gridded data set was converted to a coloured map display using the "griddata" and "meshgrid" functions and projected using an equidistant conic projection system. Vector files of the North American coastline, rivers, and lakes, and geotiff files of North American topography were imported from the public domain database Natural Earth (naturalearthdata.com).

3. Results and discussion

3.1. Isotope composition of Mojave Desert dust hotspots

The Mojave Desert is an important dust-producing region that is under-characterized geochemically ([Urban et al., 2018](#)). Our new Nd, Sr and Pb data for Mojave Desert playa lake silts ([Fig. 2](#)) improve upon existing datasets, particularly by providing a Pb isotopic characterisation. The Mojave Desert playa lake silt samples that we analysed display a modest range in Pb isotope compositions of 18.722–19.362 for ²⁰⁶Pb/²⁰⁴Pb; internal errors are ≤ ± 0.0023. ϵ_{Nd} values range from −7.5 to −12.0 (internal errors are ≤ ± 0.13 epsilon units) with the bulk of the distribution falling between −8.7 and −10.4 and relatively non-radiogenic ⁸⁷Sr/⁸⁶Sr ratios, ranging from 0.709447 to 0.716889; internal errors are ≤ ± 0.000010.

3.1.1. Mojave playa lake silts versus bedrock signatures

Radiogenic isotopes are widely used to determine the provenance of terrigenous material deposited in marine sediments by comparing the isotope composition of the deposited sedimentary material to the continental bedrock and surface sediments in potential source regions upwind (e.g. [Abouchami et al., 2013](#); [Bailey et al., 2012](#); [Lang et al., 2014](#)). This approach commonly necessitates averaging bedrock compositions over large areas of heterogeneous geology (e.g. [Lang et al., 2014](#); [Walter et al., 2000](#)). Our radiogenic maps allow us to test the fidelity of this approach. The geomorphology of the Mojave Desert is varied, both in its landforms and the geology of the country rocks ([Baldrige, 2004](#); [Jennings and Strand, 1981](#)). Aeolian material from the Mojave Desert can therefore originate from a wide array of country rock

lithologies. In Fig. 3, we compare our bedrock isotope maps to our measurements of the Mojave playa lake silts to assess similarities between proximal bedrock composition and aeolian material originating from the NASW. For all three isotope systems, our Mojave playa lake data fall within the range of the geological bedrock of the NASW region – for example, the $^{206}\text{Pb}/^{204}\text{Pb}$ isotopic composition of the geological bedrock ranges from approximately 17 to 20.5, which encompasses the spread of the measured $^{206}\text{Pb}/^{204}\text{Pb}$ values of the Mojave playa lake silts (~18.7–19.4). Some of the playas have very similar values to the local bedrock (Fig. 3a–c), particularly in Pb and Nd space. The Sr isotope map (Fig. 3c), however, clearly demonstrates that our samples from the playa lakes of the Mojave Desert are typically more radiogenic in $^{87}\text{Sr}/^{86}\text{Sr}$ than their underlying bedrock. This offset is of the correct sign to signal incongruent behaviour of rubidium (Rb) and Sr during weathering because that process typically gives rise to more radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ values in weathered sediments than in their parent bedrock (Dasch, 1969).

In summary, our Mojave Desert playa lake samples display a relatively narrow range in Pb, Nd and Sr isotopic composition when compared with the bedrock. The geochemistry and mineralogy of modern dust deposition in the Mojave indicates a well-mixed source of material (Reynolds et al., 2006), and the isotope characteristics of our playa lake samples are likely a function of this wide-scale mixing process across the desert. Next, we use these Results to understand better the origin of the silt mantles on the Californian Channel Islands.

3.2. Origin of Channel Island silt mantles, offshore Southern California

3.2.1. Assessing the likelihood of a significant local (island) source

Satellite imagery provides strong evidence for the transport of aeolian material to the Channel Islands from the NASW (Fig. 1b) but the importance of aeolian supply versus a local contribution from the island bedrock to the silt mantles must be carefully assessed. Contrasting bedrock lithology between the NASW and Channel Islands makes this a straightforward task. For the <63 μm fraction of the Channel Island silt mantles (from San Clemente and Anacapa islands) ϵ_{Nd} values range from –12.1 to –8.1 with internal errors $\leq \pm 0.12$ and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios are relatively non-radiogenic, ranging from 0.708610 to 0.710575 with internal errors $\leq \pm 0.000008$ (Fig. 2a). The samples are closely grouped in Pb isotope space, from 18.809 to 19.098 in $^{206}\text{Pb}/^{204}\text{Pb}$, from 15.644 to 15.677 in $^{207}\text{Pb}/^{204}\text{Pb}$ and from 39.051 to 39.337 in $^{208}\text{Pb}/^{204}\text{Pb}$ space (internal errors $\leq \pm 0.0063$, 0.0058 and 0.0186, respectively; Fig. 2e–h). The subset of Channel Island silt mantle samples that are <10 μm have similar values in Pb isotope space to the <63 μm fraction (Fig. 2e–h). Their Nd isotopic values have a smaller range compared to the <63 μm fraction and their average value is shifted to slightly more radiogenic (less negative) ϵ_{Nd} values (–10.3 to –8.1, internal errors $\leq \pm 0.07$, Fig. 2a). The measured $^{87}\text{Sr}/^{86}\text{Sr}$ values (0.709194–0.710109, internal errors $\leq \pm 0.000007$, Fig. 2a) of the <10 μm fraction fall within the same range as the <63 μm fraction.

The isotopic values of these mantles are strikingly different to those of Channel Island bedrock (Fig. 2a). Whole-rock measurements of $^{87}\text{Sr}/^{86}\text{Sr}$ in San Clemente Island andesites fall between 0.7037 and 0.7045 (Johnson and O'Neil, 1984). Although no bedrock Sr isotopic data are available for Anacapa Island, these andesitic bedrocks are compositionally similar and stratigraphically correlative to the volcanic suite on Santa Cruz Island (Weigand, 1993) which has $^{87}\text{Sr}/^{86}\text{Sr}$ ratios from 0.7025 to 0.7032 (Johnson and O'Neil, 1984). Similarly, Nd isotope data measured on the volcanics from the other Channel Islands (+3.4 to +9.4; Weigand and Savage (1999) and references therein) serve as a good guide for

the base rock composition on Anacapa and San Clemente. Our data (Fig. 2a) for the silt mantles contrast sharply with these island bedrock compositions (and even stream bed samples from Santa Rosa Island, Fig. 2a, Napier et al. (2020)), providing strong support for the interpretations of Muhs et al. (2007) and Muhs et al. (2008) that the island silt mantles are aeolian in origin.

3.2.2. No significant input to Channel Island silts from Asia or Africa

Next, we consider potential external sources of the Channel Island wind-blown silts. We consider three main sources for this aeolian material: (i) the distal influence of the Pacific Asian dust plume bringing material predominantly originating from East Asia (Creamean et al., 2014; Uno et al., 2011; VanCuren and Cahill, 2002), (ii) aeolian material originating from Africa, considered because the Sahara is the world's largest dust source (Ginoux et al., 2012; Prospero et al., 2002) and African dust storms have been recorded travelling as far west as Colorado and Texas (Hand et al., 2017; Perry et al., 1997), or in very rare cases, eastwards across Asia and the Pacific to the western coast of North America (McKendry et al., 2007), and (iii) transport from North America on the Santa Ana winds (Fig. 1b; Muhs et al., 2007; Muhs et al., 2008).

Comparison of similar size fractions helps to minimise the widely acknowledged effects exerted on $^{87}\text{Sr}/^{86}\text{Sr}$ by weathering and grain size (Dasch, 1969; Feng et al., 2009), with the fine silt- and clay-sized fractions of windblown material typically carrying higher $^{87}\text{Sr}/^{86}\text{Sr}$ ratios than their coarser fraction counterparts because of the preferential incorporation of K- and hence Rb-rich clay and mica minerals into the finest fractions (Feng et al., 2009). Although previous studies have shown that aeolian material transported over thousands of kilometres is borne largely in the <5 μm fraction (and often even in the <2 μm fraction (Pye, 1987a)), it is possible for larger dust particles to travel considerable distances from their source region (e.g. Betzer et al., 1988; van der Does et al., 2018a) and the Channel Islands are <150 km from potential mainland US dust sources. Therefore, we consider data for the total suspended sediment, loess and where available, we include data measured on the <75 μm fraction. Taking this approach also accounts for the break of up larger-sized dust particles during saltation and other near-surface processes into finer size fractions and their subsequent long-range transport.

To evaluate the likelihood of an Asian influence we compare our data to measurements of Chinese loess deposits, which are representative of Asian-sourced dusts (Chen et al., 2007; Jones et al., 2000; Kanayama et al., 2005; Sun and Zhu, 2010; Wu et al., 2011; Zeng et al., 2015; Zhao et al., 2015) in Fig. 2. We find that the Pb isotope composition of Asian material (Jones et al., 2000; Wu et al., 2011; Zeng et al., 2015) is distinct from material originating on the North American continent (Fig. 2e–h). The windblown island silt mantles and active North American dust sources cluster along a single, well-defined trend in $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{207}\text{Pb}/^{204}\text{Pb}$ space (Fig. 2e and f) that is offset from the distribution of the Asian data field (Fig. 2e and f). There is also minimal overlap in Nd–Sr space between the Asian data and the different sources of terrigenous material from the NASW (Fig. 2a–c). Asian loess has a broader distribution in ϵ_{Nd} space (approx. –14 to –4) and typically shows much more radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ ratios than the Channel Island silts. In fact, the subset of Channel Island silt samples that were sieved to <10 μm do not overlap with the Asian loess in Nd–Sr space at all (Fig. 2c). Thus, while Pb isotope data do not rule out the presence of Asian dust in the silt mantles when considered in isolation, when considered together with the Nd and Sr systems, radiogenic isotopes provide a powerful tool to distinguish more clearly between long-range transported Asian dust and North American-derived material (windblown and riverine) in Pacific sediments. Our data therefore clearly rule out the possibility of a major contribution




Fig. 2. Radiogenic isotope characterisation of sources and sinks of windblown material from the NASW and nearshore eastern Pacific Ocean in: (a–d) Nd–Sr space, (e–f) $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{207}\text{Pb}/^{204}\text{Pb}$ space and (g–h) $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{208}\text{Pb}/^{204}\text{Pb}$ spaces. Panel c) shows a close-up of the Nd–Sr region outlined in a) and panels b) and d) display histograms of the distribution of the ϵ_{Nd} isotope data from ODP sites 893 and 1015, distinguishing between 'riverine' and hemipelagic samples. Panel f) shows a close-up of the $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{207}\text{Pb}/^{204}\text{Pb}$ region outlined in e). Panel h) shows a close-up of the $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{208}\text{Pb}/^{204}\text{Pb}$ region outlined in g). Data for the NASW (in blue) and Transverse Ranges (TR) river sediment (in purple) (our data, [Aarons et al., 2017](#); [Napier et al., 2020](#)) are compared to data from Chinese loess deposits, as representative of Asian dust sources ([Chen et al., 2007](#); [Jones et al., 2000](#); [Kanayama et al., 2005](#); [Sun and Zhu, 2010](#); [Wu et al., 2011](#); [Zeng et al., 2015](#); [Zhao et al., 2015](#), in green) and trans-Atlantic African dust ([Bozlaker et al., 2018](#); [Kumar et al., 2014](#); [Kumar et al., 2018](#); [Meyer et al., 2011](#); [Pourmand et al., 2014](#); [Skonieczny et al., 2013](#); [Skonieczny et al., 2014](#); [van der Does et al., 2011a](#); [van der Does et al., 2018b](#), in orange). Data for the Channel Island silt mantles (this study), material from the California borderland basins from ODP sites 893 and 1015 and Pacific core data (this study, [Napier et al., 2020](#); [Rosenbauer et al., 2013](#); [Stancin et al., 2006](#)) indicate the isotopic composition of aeolian, riverine and hemipelagic sediments sourced from North America. Nd–Sr fields of the Channel Islands

from Asia to the windblown dust deposits accumulating on the California Channel Islands.

We next consider the likelihood of a significant deposition of trans-Atlantic African aeolian material on the California Channel Islands by comparing our Channel Island silt mantle data to dust trap data from North Africa, the North Atlantic and the Caribbean (Bozlaker et al., 2018; Kumar et al., 2014; Kumar et al., 2018; Meyer et al., 2011; Pourmand et al., 2014; Skonieczny et al., 2013; Skonieczny et al., 2011b; van der Does et al., 2018b, Fig. 2). While there is some overlap in Pb isotope space between our Channel Island data and the Saharan isotope fields (Fig. 2e–h), there is little overlap in Sr–Nd isotope space. We also note that there is a poor isotopic match between the Channel Islands silt mantles and the two major North African regions implicated in trans-Atlantic dust transport: the western ($^{87}\text{Sr}/^{86}\text{Sr} = 0.7279 \pm 0.0052$, $\epsilon_{\text{Nd}} = -14.8 \pm 2.2$) and central ($^{87}\text{Sr}/^{86}\text{Sr} = 0.7186 \pm 0.0053$, $\epsilon_{\text{Nd}} = -10.0 \pm 3.9$) preferential source areas (Jewell et al., 2021). Therefore, we consider it unlikely that the Saharan dust plume makes a significant contribution to sediments offshore Southern California.

3.2.3. North American sources of Channel Island aeolian silt mantles

Next, we evaluate possible North American sources of dust. Both the Mojave Desert and the coastal mountains of Southern California are proposed as potential source regions of aeolian supply to the Channel Islands (Muhs et al., 2007, 2008). Both of these regions contain playa lakes and river valleys, which are known sources of readily deflated silt-sized material (Pye, 1987b). We compare the Sr–Nd–Pb isotope composition of the Channel Island silt mantles to our Mojave playa lake silt data and other published records (Aarons et al., 2017; Napier et al., 2020) in Fig. 2.

The <63 μm fraction of the Channel Island silt mantle samples shows a very similar trend in $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{207}\text{Pb}/^{204}\text{Pb}$ space to that of the Mojave playa lake silts (Fig. 2e). In $^{206}\text{Pb}/^{204}\text{Pb}$ vs $^{208}\text{Pb}/^{204}\text{Pb}$ space, the <63 μm fraction of the Channel Island silt mantles are completely contained within the spread of the Mojave playa lake data (Fig. 2g). Thus, on the basis of our Pb isotope data, the Mojave Desert is a compatible source for the aeolian material mantling the Channel Islands. This conclusion is supported by the Nd isotope data because the composition of the Channel Island silt mantles is very similar to the Mojave Desert samples (Fig. 2a,c). There is a slight discrepancy in Sr values between the Channel Island silts and the Mojave Desert playa lake silts (Fig. 2c) but the Channel Island data fall within the spread of dust sources from the Mojave (Fig. 2).

A geochemical characterization of the coastal mountains of southern California is provided by Napier et al. (2020) who sampled streambed sediments in the coastal California Transverse ranges (Fig. 2a,c). There is extensive overlap in Sr–Nd space between the field described by those data and our data for the Mojave playa lakes and the Channel Island silt mantles (Fig. 2). A study of potential dust sources across a wider area of the North American West shows that several other regions including the Colorado Plateau and Basin and Range province also show similar radiogenic isotope fingerprints to the Mojave (Aarons et al., 2017). Therefore, we cannot precisely identify the contribution to the island deposits made by each of these different source regions relative to the Mojave Desert. Regardless, our data strongly support the suggestion that these deposits are derived from North America (Muhs

et al., 2007, 2008) and the pathway of the Santa Ana winds across Southern California to the offshore Pacific (Conil and Hall, 2006; Hughes and Hall, 2010) suggests that active dust sources along this route, including those of the Mojave, are likely to be the most important contributors to the Channel Island silt mantles and surrounding ocean sediments.

3.3. Distinguishing sources of terrigenous deposition offshore Southern California

The Channel Island silt mantles provide an excellent case study to characterize the signature of aeolian material exported to the Eastern Pacific, but to find readily datable continuous palaeoclimate archives spanning many thousands or millions of years, we must turn instead to marine deposits that can contain terrigenous inputs derived via both riverine and aeolian transport.

Previous studies investigating sediment delivery to the California borderland basins concluded that fluvial input dominates the sediment budget (>80% in some parts of the inner borderlands, location Fig. 1c); whereas biogenic material contributes typically 20–25% (Inman and Jenkins, 1999; Schwabach and Gorsline, 1985). However, more recent work points to a significant additional input of aeolian material (Muhs et al., 2007). Although the predominant transport direction of dust from the NASW is eastwards on westerly winds towards the North American interior and possibly the distant Atlantic Ocean (Mahowald et al., 2006), at certain times during autumn and winter, ‘Santa Ana’ winds carry dust westwards to the Pacific Ocean ((Muhs et al., 2007; Muhs et al., 2008; Reheis et al., 1995) Fig. 1b). It is estimated that aeolian materials contribute anywhere between 5 and 60% of the sediment deposited in the California Borderlands, which varies primarily as a function of dilution by riverine material and biogenic inputs (Muhs et al., 2007). Here, we explore whether radiogenic isotopes can be used to assess the relative importance of aeolian and fluvial inputs to the hemipelagic sediments at ODP sites 893 and 1015.

To isolate the signature of the riverine inputs exported offshore Southern California, we analysed samples from marine drill cores at ODP sites 893 and 1015 containing fluvial deposit horizons (the ODP Site 893 ‘grey-layers’ and ODP Site 1015 turbidites). We compare the radiogenic isotope composition of hemipelagic sediments from ODP Site 1015 (which should contain both aeolian and riverine material) to these fluvially derived horizons and to the Channel Island silt mantles to understand better the relative importance of these two sources to hemipelagite formation in the region. The subset of our Californian margin borderland basin samples (located in the Santa Barbara and Santa Monica basins (Fig. 1c)) which are dominantly fluvially derived have $^{87}\text{Sr}/^{86}\text{Sr}$ values between 0.710973 and 0.715886 (internal errors $\leq \pm 0.000007$) and ϵ_{Nd} values between –13.0 and –11.4 (internal errors $\leq \pm 0.14$). The isotopic signatures of the fluvially derived layers in the Santa Barbara (ODP Site 893) and Santa Monica (ODP Site 1015) basins are very similar, with the samples from ODP Site 1015 recording slightly more unradiogenic ϵ_{Nd} and lower $^{206}\text{Pb}/^{204}\text{Pb}$ values (Fig. 2) than ODP Site 893. Holocene and deglacial samples from ODP Site 1015 do not show an appreciable compositional difference attributable to age.

In Pb isotope space, the hemipelagic data are not offset from the fluvially sourced samples (Fig. 2e–h), but Nd–Sr measurements of the ODP core samples (Fig. 2a,c) indicate a subtle but quantifiable isotopic difference between horizons characterised as fluvial and

bedrock (Johnson and O’Neil, 1984; Weigand, 1993; Weigand and Savage, 1999) and stream bed material from Santa Rosa Island (Napier et al., 2020) are also shown in panel a). Key abbreviations: CISM = Channel Island silt mantles; SBB = Santa Barbara Basin. Northern Hemisphere Regression Line (NHRL) is shown on Pb isotope plots for reference (Hart, 1984). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




Fig. 3. Radiogenic isotope maps of the bedrock geology of the North American Southwest (NASW) in interpolated a) $^{206}\text{Pb}/^{204}\text{Pb}$; b) ϵ_{Nd} ; c) $^{87}\text{Sr}/^{86}\text{Sr}$ space compared to data for unconsolidated sediment samples (this study) from Mojave Playa Lake silts, Channel Island Silt Mantles, DSDP and ODP sites (coloured circles). Colour scale (right) applies to both bedrock geology and sediment samples. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




Fig. 4. Radiogenic isotope characterisation of sources of Asian, South/Central American and North American aeolian material compared to the radiogenic isotope signature recorded in eastern Pacific marine sediment cores downwind of these continents in a) $^{208}\text{Pb}/^{206}\text{Pb}$ vs $^{207}\text{Pb}/^{206}\text{Pb}$ space and b) Nd-Sr space. Panel a) displays Pb isotope ratios plotted as $^{208}\text{Pb}/^{206}\text{Pb}$ vs $^{207}\text{Pb}/^{206}\text{Pb}$ to be consistent with Stancin et al. (2006). The sediment cores are grouped into three fields representing the dominant source as identified by Stancin et al. (2006). We have updated these three isotopic fields to include new data from both the original sites and new ones. c) Locations of the Pacific marine sediment cores used to characterize the radiogenic isotope fields displayed in panels a) and b). Colours of the markers correspond to the interpreted dominant source of terrigenous material at that core site. Red = Asia; blue = North America; purple = South/Central America. The marker shape indicates whether the data generated from that sediment core were included in the analysis of Stancin et al. (2006). Diamonds = sediment cores for which we use the same data as Stancin et al. (2006); triangles = sediment cores used by Stancin et al. (2006) with additional data incorporated here; circles = sediment cores not included in Stancin et al. (2006). Note that Sr isotope data are not included from ODP sites 848-851 due to the influence of marine barite on sediment signatures at these sites (Xie and Marcantonio, 2012). Data sources are listed in the Supplementary Information. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.).

'hemipelagic background' (Kennett et al., 1994; Lyle et al., 1997). The ϵ_{Nd} values of the hemipelagic material (which is expected to contain both aeolian and fluvial inputs) are more radiogenic (-9.4 to -9.2 ; internal errors $\leq \pm 0.08$) than the fluvial samples (-13.0 to -11.4) and the $^{87}\text{Sr}/^{86}\text{Sr}$ composition plots at the less radiogenic end of the range in the riverine samples (Fig. 2b–d).

The distinct isotopic composition of the hemipelagic sediments from ODP Site 1015 implies either a shift in the source of terrigenous material deposited in the absence of large fluvial inputs to the Santa Monica Basin or a mixing of the fluvial inputs with a second, isotopically distinct source of terrigenous material. Variability in the signature of the riverine inputs is supported by Napier et al. (2020), who attribute more radiogenic ϵ_{Nd} and less radiogenic $^{87}\text{Sr}/^{86}\text{Sr}$ values in sediment samples from the Santa Barbara Basin of Last Glacial Maximum age to an increased contribution from the southern slopes of the Santa Ynez and Topatopa Mountains. Alternatively, we note that hemipelagic sediment of both Holocene and deglacial age from ODP Site 1015 in the Santa Monica Basin more closely resemble the Mojave Desert and Channel Island silt mantles than the dominantly fluvial material sampled at the same site in Nd–Sr space, supporting a greater influence of windblown dust from the NASW during times of hemipelagic deposition (Fig. 2a).

In summary, we identify differences in Sr–Nd isotopic signatures between river-derived and hemipelagic sediments immediately offshore Southern California possibly attributable to differences in the geochemical signatures of aeolian and riverine sediments to the Californian margin. However, overlap between the isotopic signatures of terrestrial samples of source regions in the Mojave, Transverse Range and greater NASW mean that it is not yet possible to estimate the relative proportions of riverine and aeolian inputs to the Inner California Borderland Basins using Sr–Nd–Pb isotopes (Fig. 2). We therefore suggest that the Outer California Borderland Basins are a more attractive proposition for studying past dust flux and palaeoaridity in the North American Southwest because inputs of riverine material there are much lower than to the Inner Borderland Basins (Schwalbach and Gorsline, 1985).

3.4. Spatial variations in the geochemistry of hemipelagic deposition in the Eastern Pacific

In their study of the provenance of terrigenous input to the Pacific Ocean, Stancin et al. (2006) analysed the lithogenic fraction from drill cores across the eastern Pacific to determine the relative contributions from Asia, South/Central America and North America. They identified two sediment core sites as containing a terrigenous fraction dominantly sourced from North America: Deep Sea Drilling Project (DSDP) sites 32 and 469. DSDP Site 469 is located at 32.61°N , south of ODP sites 893 and 1015, but it is further offshore than either of these two ODP sites, situated at the base of the Patton Escarpment bounding the Southern Californian Outer Borderland Basin (Figs. 1 and 3). In contrast, DSDP Site 32 is located at 37.12°N , a similar latitude to San Francisco Bay, approximately 820 km north of DSDP Site 469 (Fig. 3). Radiogenic isotope data of Plio–Pleistocene samples from these two sites were used to determine the North American end member in Pb, Sr and Nd space for the entirety of the North Pacific Ocean (Stancin et al., 2006). Our data provide a test of that approach.

3.4.1. North American terrigenous supply to the North Pacific Ocean

DSDP sites 32 and 469 are isotopically distinct from each other in Nd–Sr and Pb isotope space (Fig. 2). DSDP Site 32 is located (Fig. 3) proximal to the Delgada Submarine Fan (McManus et al., 1970), which consists of terrigenous material derived mainly from the San Joaquin and Sacramento river catchments and our Pb

and Sr isotope maps show good isotopic agreement between the drainage basins of these two rivers (Fig. 3, see also Supplementary Fig. 2) and DSDP Site 32. Furthermore, there is good isotopic correspondence between the DSDP Site 32 data (Stancin et al., 2006) and sedimentary grab samples from the mouth and near offshore of San Francisco Bay where the San Joaquin and Sacramento Rivers enter the Pacific Ocean (Fig. 2a, data labelled as 'Northern California offshore sediments' (Rosenbauer et al., 2013)). Taken together, these lines of evidence strongly suggest that the dominant source of the hemipelagic deposition at DSDP Site 32 is derived from fluvial supply from Northern California.

DSDP Site 469 is located well to the south of DSDP Site 32, just seaward of the Southern Californian Borderland basin system and the Channel Islands that lie therein (Figs. 1 and 3). Here, the influence of Northern Californian rivers is suggested to be minimal (Griggs and Hein, 1980) and our analysis substantiates this suggestion because, in contrast to DSDP Site 32, the terrigenous fraction from DSDP Site 469 shows significant isotopic correspondence with Channel Island silts and NASW dust sources (Fig. 2, this study, Aarons et al. (2017); Stancin et al. (2006)). These findings suggest that aeolian-sourced material from the NASW is a significant contributor to hemipelagic deposition at DSDP Site 469.

In Fig. 4 we compare the isotopic composition of the terrigenous fraction in Pacific Ocean marine sediment cores with data from continental sediment sources, including our new data for North America. The comparison shows much greater ranges in isotopic composition of the sediment sources than the sediment sinks (the cores), highlighting the extent of signal homogenisation in the marine core data caused by mixing of contributions during transport over long distances.

Our comparisons clearly highlight the advantage of considering multiple isotope systems to characterise and trace long-range aeolian transport and deposition. For example, there is considerable overlap in Nd–Sr isotope space between North American sediment sources and the South/Central American-dominated provenance grouping in the sediment core data (Fig. 4b) but a major North American influence on the eastern equatorial Pacific can be ruled out by the clear distinction in Pb isotope space illustrated in Fig. 4a.

More work is needed to pinpoint the contributions of different dust sources in western North America but our data provide a solid framework to help trace the extent and temporal variability of North American dust deposition in the Pacific Ocean. Current data coverage in the Northeast Pacific is sparse, with sites from both riverine and dust-dominated locations included in Fig. 4. Detailed characterisation of the aeolian sediment source region signatures permits a more accurate assessment of their long-range contributions than can be provided by marine sediments where riverine and aeolian inputs from multiple sources commonly mix. Our results support the use of radiogenic isotopes to discriminate between Asian and North American sourced dusts in Pacific Ocean sediments. However, distinguishing aeolian inputs from riverine supply along the Californian margin using radiogenic isotopes alone is not straightforward. We suggest that future work on dust export from the NASW should focus on more sites in the Outer California Borderland Basins and eastern Pacific margin where the influence of riverine supply is smaller than for the Inner Borderland Basins.

4. Conclusions

We present new radiogenic isotope data to characterise sources and sinks of terrigenous material originating from the North American Southwest in conjunction with new Pb, Sr and Nd isotope maps of the geological bedrock of the region capturing the broad-scale features of bedrock that help to trace sources and sinks of

aeolian material. Our radiogenic isotope data improve our ability to geochemically fingerprint terrigenous material (both aeolian and fluvial) from the North American Southwest (NASW) and permit clear distinction between windblown material from the NASW and long-range transported material from Asia and North Africa. We rule out a significant contribution from either Asia or North Africa to the Quaternary silts mantling the California Channel Islands and corroborate the suggestion that they are aeolian in origin and sourced from the NASW on the Santa Ana winds.

We conclude that aeolian-sourced material from the NASW is a significant contributor to marine sediments offshore Southern California. The geochemical distinction between riverine and hemipelagic deposits reveals the potential of the sediments of the Outer California Borderland Basins to preserve a signature of past aeolian inputs and hence produce high-fidelity reconstructions of dust export and palaeoaridity in the NASW over geological timescales.

Credit author statement

Jardine, G.E.: Conceptualisation, Methodology, Investigation, Formal analysis, Writing – original draft, Writing – review & editing, Visualisation, Crocker A.J.: Conceptualisation, Formal analysis, Writing – original draft, Writing – review & editing, Supervision, Bailey, I.: Conceptualisation, Writing – original draft, Writing – review & editing, Supervision, Cooper, M.J.: Methodology, Investigation, Resources, Validation, Writing – review & editing, Milton, J.A.: Methodology, Investigation, Resources, Validation, Wilson, P.A.: Conceptualisation, Formal analysis, Writing – original draft, Writing – review & editing, Supervision.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank D. Muhs at the USGS for samples and valuable insights during the course of this research. Sediment core samples were provided by the IODP, funded by the U.S. National Science Foundation and participating countries. We are also grateful to R. Taylor, A. Michalik and C. Bristow for laboratory assistance and constructive discussions and to three anonymous reviewers whose comments greatly improved this manuscript. This study was supported by the University of Southampton in the form of a Mayflower Scholarship to G.E.J. I.B. acknowledges funding from an EU MARINEFF project (Interreg VA France-(Channel)-England Programme project #162). P.A.W. and A.J.C. acknowledge Royal Society funding including a Wolfson Merit Award to P.A.W.

Data availability

The data presented in this manuscript are available as a supplementary data table and will also be deposited in the PANGAEA open access library.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.quascirev.2021.106934>.

References

- Aarons, S.M., Blakowski, M.A., Aciego, S.M., Stevenson, E.I., Sims, K.W.W., Scott, S.R., Aarons, C., 2017. Geochemical characterization of critical dust source regions in the American West. *Geochim. Cosmochim. Acta* 215, 141–161.
- Abouchami, W., N  the, K., Kumar, A., Galer, S.J.G., Jochum, K.P., Williams, E., Horbe, A.M.C., Rosa, J.W.C., Balsam, W., Adams, D., Mezger, K., Andreae, M.O., 2013. Geochemical and isotopic characterization of the Bod  le Depression dust source and implications for transatlantic dust transport to the Amazon Basin. *Earth Planet. Sci. Lett.* 380, 112–123.
- Abouchami, W., Zabel, M., 2003. Climate forcing of the Pb isotope record of terrigenous input into the Equatorial Atlantic. *Earth Planet. Sci. Lett.* 213, 221–234.
- Ahn, J., Brook, E.J., Mitchell, L., Rosen, J., McConnell, J.R., Taylor, K., Etheridge, D., Rubino, M., 2012. Atmospheric CO₂ over the last 1000 years: a high-resolution record from the west Antarctic ice sheet (WAIS) divide ice core. *Global Biogeochem. Cycles* 26.
- Ault, T.R., Mankin, J.S., Cook, B.I., Smerdon, J.E., 2016. Relative impacts of mitigation, temperature, and precipitation on 21st-century megadrought risk in the American Southwest. *Sci. Adv.* 2.
- Bailey, I., Foster, G.L., Wilson, P.A., Jovane, L., Storey, C.D., Trueman, C.N., Becker, J., 2012. Flux and provenance of ice-rafted debris in the earliest Pleistocene sub-polar North Atlantic Ocean comparable to the last glacial maximum. *Earth Planet. Sci. Lett.* 341–344, 222–233.
- Baldrige, W.S., 2004. *Geology of the American Southwest: A Journey through Two Billion Years of Plate-Tectonic History*. Cambridge University Press, United Kingdom.
- Balling, R.C., Goodrich, G.B., 2010. Increasing drought IN the AMERICAN southwest? A continental perspective using a spatial analytical evaluation OF recent trends. *Phys. Geogr.* 31, 293–306.
- Bayon, G., German, C.R., Boella, R.M., Milton, J.A., Taylor, R.N., Nesbitt, R.W., 2002. An improved method for extracting marine sediment fractions and its application to Sr and Nd isotopic analysis. *Chem. Geol.* 187, 179–199.
- Behl, R.J., 1995. Sedimentary facies and sedimentology of the late quaternary Santa Barbara Basin, site 893. *Proceedings of the ocean drilling Program. Sci. Results* 146, 295–308.
- Betzler, P.R., Carder, K.L., Duce, R.A., Merrill, J.T., Tindale, N.W., Uematsu, M., Costello, D.K., Young, R.W., Feely, R.A., Breland, J.A., Bernstein, R.E., Greco, A.M., 1988. Long-range transport of giant mineral aerosol particles. *Nature* 336, 568–571.
- Bozlaker, A., Prospero, J.M., Price, J., Chellam, S., 2018. Linking Barbados mineral dust aerosols to North African sources using elemental composition and radiogenic Sr, Nd, and Pb isotope signatures. *J. Geophys. Res.: Atmospheres* 123, 1384–1400.
- Cayan, D.R., Das, T., Pierce, D.W., Barnett, T.P., Tyree, M., Gershunov, A., 2010. Future dryness in the southwest US and the hydrology of the early 21st century drought. *Proc. Natl. Acad. Sci. U. S. A.* 107, 21271–21276.
- Chen, J., Li, G.J., Yang, J.D., Rao, W.B., Lu, H.Y., Balsam, W., Sun, Y.B., Ji, J.F., 2007. Nd and Sr isotopic characteristics of Chinese deserts: implications for the provenances of Asian dust. *Geochim. Cosmochim. Acta* 71, 3904–3914.
- Cole, J.M., Goldstein, S.L., Demenocal, P.B., Hemming, S.R., Grousset, F.E., 2009. Contrasting compositions of Saharan dust in the eastern Atlantic Ocean during the last deglaciation and African humid period. *Earth Planet. Sci. Lett.* 278, 257–266.
- Conil, S., Hall, A., 2006. Local regimes of atmospheric variability: a case study of southern California. *J. Clim.* 19, 4308–4325.
- Cook, B.I., Ault, T.R., Smerdon, J.E., 2015. Unprecedented 21st century drought risk in the American southwest and central plains. *Sci. Adv.* 1, e1400082.
- Cook, B.I., Smerdon, J.E., Seager, R., Cook, E.R., 2014. Pan-continental droughts in north America over the last millennium. *J. Clim.* 27, 383–397.
- Cook, E.R., Seager, R., Cane, M.A., Stahle, D.W., 2007. North American drought: reconstructions, causes, and consequences. *Earth Sci. Rev.* 81, 93–134.
- Creamean, J.M., Spackman, J.R., Davis, S.M., White, A.B., 2014. Climatology of long-range transported Asian dust along the west coast of the United States. *J. Geophys. Res.: Atmospheres* 119, 2014JD021694.
- Dasch, E.J., 1969. Strontium isotopes in weathering profiles, deep-sea sediments, and sedimentary rocks. *Geochim. Cosmochim. Acta* 33, 1521–1552.
- de la Vega, E., Chalk, T.B., Wilson, P.A., Bysani, R.P., Foster, G.L., 2020. Atmospheric CO₂ during the mid-piacenzian warm period and the M2 glaciation. *Sci. Rep.* 10, 11002.
- Feakins, S.J., Kirby, M.E., Cheetham, M.I., Ibarra, Y., Zimmerman, S.R.H., 2014. Fluctuation in leaf wax D/H ratio from a southern California lake records significant variability in isotopes in precipitation during the late Holocene. *Org. Geochem.* 66, 48–59.
- Feng, J.L., Zhu, L.P., Zhen, X.L., Hu, Z.G., 2009. Grain size effect on Sr and Nd isotopic compositions in eolian dust: implications for tracing dust provenance and Nd model age. *Geochim. J.* 43, 123–131.
- Ginoux, P., Prospero, J.M., Gill, T.E., Hsu, N.C., Zhao, M., 2012. Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products. *Rev. Geophys.* 50, RG3005.
- Griggs, G.B., Hein, J.R., 1980. Sources, dispersal, and clay mineral composition of fine-grained sediment off central and northern California. *J. Geol.* 88, 541–566.
- Hand, J.L., Gill, T.E., Schichtel, B.A., 2017. Spatial and seasonal variability in fine mineral dust and coarse aerosol mass at remote sites across the United States.

- J. Geophys. Res.: Atmospheres 122, 3080–3097.
- Hart, S.R., 1984. A large-scale isotope anomaly in the Southern Hemisphere mantle. *Nature* 309, 753–757.
- Hoell, A., Funk, C., Barlow, M., 2014. The regional forcing of Northern hemisphere drought during recent warm tropical west Pacific Ocean La Nina events. *Clim. Dynam.* 42, 3289–3311.
- Hughes, M., Hall, A., 2010. Local and synoptic mechanisms causing Southern California's Santa Ana winds. *Clim. Dynam.* 34, 847–857.
- Hyeong, K., Kim, J., Pettke, T., Yoo, C.M., Hur, S.-d., 2011. Lead, Nd and Sr isotope records of pelagic dust: source indication versus the effects of dust extraction procedures and authigenic mineral growth. *Chem. Geol.* 286, 240–251.
- Inman, D.L., Jenkins, S.A., 1999. Climate change and the episodicity of sediment flux of small California rivers. *J. Geol.* 107, 251–270.
- Jacobsen, S.B., Wasserburg, G.J., 1980. SM-ND isotopic evolution OF chondrites. *Earth Planet Sci. Lett.* 50, 139–155.
- Jennings, C.W., Strand, R.G., 1981. Geologic Map of California, fifth ed.
- Jewell, A.M., Drake, N., Crocker, A.J., Bakker, N.L., Kunkelova, T., Bristow, C.S., Cooper, M.J., Milton, J.A., Breeze, P.S., Wilson, P.A., 2021. Three North African dust source areas and their geochemical fingerprint. *Earth Planet Sci. Lett.* 116645.
- Johnson, C.M., O'Neil, J.R., 1984. Triple junction magmatism: a geochemical study of Neogene volcanic rocks in western California. *Earth Planet Sci. Lett.* 71, 241–262.
- Jones, C.E., Halliday, A.N., Rea, D.K., Owen, R.M., 2000. Eolian inputs of lead to the North Pacific. *Geochim. Cosmochim. Acta* 64, 1405–1416.
- Just, J., Heslop, D., von Döbenek, T., Bickert, T., Dekkers, M.J., Frederichs, T., Meyer, I., Zabel, M., 2012. Multiproxy characterization and budgeting of terrigenous end-members at the NW African continental margin. *G-cubed* 13.
- Kanayama, S., Yabuki, S., Zeng, F., Liu, M., Shen, Z.-z., Liu, L., Yanagisawa, F., Abe, O., 2005. Size-dependent geochemical characteristics of Asian dust - Sr and Nd isotope compositions as tracers for source identification. *J. Meteorol. Soc. Jpn.* 83, 107–120.
- Keeling, C.D., Piper, S.C., Bacastow, R.B., Wahlen, M., Whorf, T.P., Heimann, M., Meijer, H.A., 2001. Exchanges of Atmospheric CO₂ and 13CO₂ with the Terrestrial Biosphere and Oceans from 1978 to 2000. *Scripps Institution of Oceanography, San Diego*.
- Kennett, J.P., Baldauf, J.G., Shipboard Scientific Party, 1994. Visual Core Description from ODP Hole 143-893B.
- Kumar, A., Abouchami, W., Galer, S.J.G., Garrison, V.H., Williams, E., Andreae, M.O., 2014. A radiogenic isotope tracer study of transatlantic dust transport from Africa to the Caribbean. *Atmos. Environ.* 82, 130–143.
- Kumar, A., Abouchami, W., Galer, S.J.G., Singh, S.P., Fomba, K.W., Prospero, J.M., Andreae, M.O., 2018. Seasonal radiogenic isotopic variability of the African dust outflow to the tropical Atlantic Ocean and across to the Caribbean. *Earth Planet Sci. Lett.* 487, 94–105.
- Lang, D.C., Bailey, I., Wilson, P.A., Beer, C.J., Bolton, C.T., Friedrich, O., Newsam, C., Spencer, M.R., Gutjahr, M., Foster, G.L., Cooper, M.J., Milton, J.A., 2014. The transition on North America from the warm humid Pliocene to the glaciated Quaternary traced by eolian dust deposition at a benchmark North Atlantic Ocean drill site. *Quat. Sci. Rev.* 93, 125–141.
- Larrasoana, J.C., Roberts, A.P., Rohling, E.J., Winkhofer, M., Wehausen, R., 2003. Three million years of monsoon variability over the northern Sahara. *Clim. Dynam.* 21, 689–698.
- Lyle, M., Koizumi, I., Richter, C., Shipboard Scientific Party, 1997. Visual Core Description from ODP Hole 167-1015B.
- Macdonald, G.M., Stahle, D.W., Diaz, J.V., Beer, N., Busby, S.J., Cerano-Paredes, J., Cole, J.E., Cook, E.R., Endfield, G., Gutierrez-Garcia, G., Hall, B., Magan, V., Meko, D.M., Méndez-Pérez, M., Sauchyn, D.J., Watson, E., Woodhouse, C.A., 2008. Climate warming and 21st-century drought in Southwestern north America. *Eos. Trans. Am. Geophys. Union* 89, 82–82.
- Mahowald, N.M., Muhs, D.R., Levis, S., Rasch, P.J., Yoshioka, M., Zender, C.S., Luo, C., 2006. Change in atmospheric mineral aerosols in response to climate: last glacial period, preindustrial, modern, and doubled carbon dioxide climates. *Journal of Geophysical Research. Atmospheres* 111 n/a-n/a.
- Marsaglia, K.M., Rimkus, K.C., Behl, R.J., 1995. Provenance of sand deposited in the Santa Barbara Basin at site 893 during the last 155,000 years. *Proceedings of the ocean drilling Program. Sci. Results* 146, 61–75.
- McKendry, I.G., Strawbridge, K.B., O'Neill, N.T., Macdonald, A.M., Liu, P.S.K., Leitch, W.R., Anlauf, K.G., Jaegle, L., Fairlie, T.D., Westphal, D.L., 2007. Trans-Pacific transport of Saharan dust to western North America: a case study. *J. Geophys. Res.: Atmospheres* 112.
- McManus, D.A., Burns, R.E., von der Borch, C.C., Goll, R.M., Milow, E.D., Olsson, R.K., Vallier, T., Weser, O., 1970. Site 32. *Affiliation (Analytic): Univ. Wash., Seattle, WA*, pp. 5–15.
- Medellín-Azuara, J., MacEwan, D., Howitt, R.E., Sumner, D.A., Lund, J.R., 2016. Economic Impacts of the 2016 California Drought for Agriculture, p. 17.
- Meinshausen, M., Smith, S.J., Calvin, K., Daniel, J.S., Kainuma, M.L.T., Lamarque, J.F., Matsumoto, K., Montzka, G.A., Raper, S.C.B., Riahi, K., Thomson, A., Velders, G.J.M., van Vuuren, D.P.P., 2011. The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Climatic Change* 109.
- Meyer, I., Davies, G.R., Stuut, J.-B.W., 2011. Grain size control on Sr-Nd isotope provenance studies and impact on paleoclimate reconstructions: an example from deep-sea sediments offshore NW Africa. *G-cubed* 12.
- Muhs, D.R., Budahn, J., Reheis, M., Beann, J., Skipp, G., Fisher, E., 2007. Airborne dust transport to the eastern Pacific Ocean off southern California: evidence from san Clemente island. *J. Geophys. Res.:h-Atmospheres* 112.
- Muhs, D.R., Budahn, J.R., Johnson, D.L., Reheis, M., Beann, J., Skipp, G., Fisher, E., Jones, J.A., 2008. Geochemical evidence for airborne dust additions to soils in Channel Islands National Park, California. *Geol. Soc. Am. Bull.* 120, 106–126.
- Nakai, S., Halliday, A.N., Rea, D.K., 1993. Provenance OF dust IN the pacific-ocean. *Earth Planet Sci. Lett.* 119, 143–157.
- Napier, T.J., Hendy, I.L., Fahnestock, M.F., Bryce, J.G., 2020. Provenance of detrital sediments in Santa Barbara Basin, California, USA: changes in source contributions between the last glacial maximum and Holocene. *GSA Bulletin* 132, 65–84.
- NASA, 2002. Dusty Skies over Southern California.
- Nizou, J., Hanebuth, T.J.J., Vogt, C., 2011. Deciphering signals of late Holocene fluvial and aeolian supply from a shelf sediment depocentre off Senegal (north-west Africa). *J. Quat. Sci.* 26, 411–421.
- Perry, K.D., Cahill, T.A., Eldred, R.A., Dutcher, D.D., Gill, T.E., 1997. Long-range transport of North African dust to the eastern United States. *J. Geophys. Res.: Atmospheres* 102, 11225–11238.
- Pettke, T., Halliday, A.N., Hall, C.M., Rea, D.K., 2000. Dust production and deposition in Asia and the North Pacific Ocean over the past 12 myr. *Earth Planet Sci. Lett.* 178, 397–413.
- Pourmand, A., Prospero, J.M., Sharifi, A., 2014. Geochemical fingerprinting of trans-Atlantic African dust based on radiogenic Sr-Nd-Hf isotopes and rare earth element anomalies. *Geology* 42, 675–678.
- Povea, P., Cacho, I., Moreno, A., Menendez, M., Mendez, F.J., 2015. A new procedure for the lithic fraction characterization in marine sediments from high productivity areas: optimization of analytical and statistical procedures. *Limnol Oceanogr. Methods* 13, 127–137.
- Prospero, J.M., Ginoux, P., Torres, O., Nicholson, S.E., Gill, T.E., 2002. Environmental characterization of global sources of atmospheric soil dust identified with the NIMBUS 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product. *Rev. Geophys.* 40, 1002.
- Pye, K., 1987a. Chapter 3-DUST ENTRAINMENT, TRANSPORT and DEPOSITION, Aeolian Dust and Dust Deposits. Academic Press, pp. 29–62.
- Pye, K., 1987b. Chapter 4-DUST SOURCES, SINKS and RATES of DEPOSITION, Aeolian Dust and Dust Deposits. Academic Press, pp. 63–91.
- Rea, D.K., 1994. The paleoclimatic record provided by eolian deposition in the deep sea: the geologic history of wind. *Rev. Geophys.* 32, 159–195.
- Reheis, M.C., Goodmacher, J.C., Harden, J.W., McFadden, L.D., Rockwell, T.K., Shroba, R.R., Sowers, J.M., Taylor, E.M., 1995. Quaternary soils and dust deposition IN southern Nevada and California. *Geol. Soc. Am. Bull.* 107, 1003–1022.
- Reynolds, R.L., Reheis, M., Yount, J., Lamothe, P., 2006. Composition of aeolian dust in natural traps on isolated surfaces of the central Mojave Desert—insights to mixing, sources, and nutrient inputs. *J. Arid Environ.* 66, 42–61.
- Rosenbauer, R.J., Foxgrover, A.C., Hein, J.R., Swarzenski, P.W., 2013. A Sr–Nd isotopic study of sand-sized sediment provenance and transport for the San Francisco Bay coastal system. *Mar. Geol.* 345, 143–153.
- Routson, C.C., Overpeck, J.T., Woodhouse, C.A., Kenney, W.F., 2016. Three millennia of Southwestern north American dustiness and future implications. *PLoS One* 11.
- Salzer, M.W., Kipfmüller, K.F., 2005. Reconstructed temperature and precipitation on a millennial timescale from tree-rings in the Southern Colorado Plateau, USA. *Climatic Change* 70, 465–487.
- Schwalbach, J.R., Gorsline, D.S., 1985. Holocene sediment budgets for the basins OF the California continental borderland. *J. Sediment. Petrol.* 55, 829–842.
- Seager, R., Ting, M., Held, I., Kushnir, Y., Lu, J., Vecchi, G., Huang, H.-P., Harnik, N., Leetmaa, A., Lau, N.-C., Li, C., Velez, J., Naik, N., 2007. Model projections of an imminent transition to a more arid climate in Southwestern north America. *Science* 316, 1181–1184.
- Seo, I., Lee, Y.I., Kim, W., Yoo, C.M., Hyeong, K., 2015. Movement of the Intertropical Convergence Zone during the mid-pleistocene transition and the response of atmospheric and surface ocean circulations in the central equatorial Pacific. *G-cubed* 16, 3973–3981.
- Seo, I., Lee, Y.I., Yoo, C.M., Kim, H.J., Hyeong, K., 2014. Sr–Nd isotope composition and clay mineral assemblages in eolian dust from the central Philippine Sea over the last 600 kyr: implications for the transport mechanism of Asian dust. *J. Geophys. Res.: Atmospheres* 119, 2014JD022025.
- Skonieczny, C., Bory, A., Bout-Roumazelles, V., Abouchami, W., Galer, S.J.G., Crosta, X., Diallo, A., Ndiaye, T., 2013. A three-year time series of mineral dust deposits on the West African margin: sedimentological and geochemical signatures and implications for interpretation of marine paleo-dust records. *Earth Planet Sci. Lett.* 364, 145–156.
- Skonieczny, C., Bory, A., Bout-Roumazelles, V., Abouchami, W., Galer, S.J.G., Crosta, X., Stuut, J.B., Meyer, I., Chiapello, I., Podvin, T., Chatenet, B., Diallo, A., Ndiaye, T., 2011a. The 7–13 March 2006 major Saharan outbreak: multiproxy characterization of mineral dust deposited on the West African margin. *Journal of Geophysical Research. Atmospheres* 116.
- Skonieczny, C., Bory, A., Bout-Roumazelles, V., Abouchami, W., Galer, S.J.G., Crosta, X., Stuut, J.B., Meyer, I., Chiapello, I., Podvin, T., Chatenet, B., Diallo, A., Ndiaye, T., 2011b. The 7–13 March 2006 major Saharan outbreak: multiproxy characterization of mineral dust deposited on the West African margin. *Journal of Geophysical Research. Atmospheres* 116.
- Stancin, A.M., Gleason, J.D., Rea, D.K., Owen, R.M., Moore, T.C., Blum, J.D., Hovan, S.A., 2006. Radiogenic isotopic mapping of late Cenozoic eolian and hemipelagic sediment distribution in the east-central Pacific. *Earth Planet Sci. Lett.* 248, 840–850.
- Sun, J.M., Zhu, X.K., 2010. Temporal variations in Pb isotopes and trace element

- concentrations within Chinese eolian deposits during the past 8 Ma: implications for provenance change. *Earth Planet Sci. Lett.* 290, 438–447.
- Super, J.R., Thomas, E., Pagani, M., Huber, M., O'Brien, C., Hull, P.M., 2018. North Atlantic temperature and pCO₂ coupling in the early-middle Miocene. *Geology* 46, 519–522.
- Tanaka, T., Togashi, S., Kamioka, H., Amakawa, H., Kagami, H., Hamamoto, T., Yuhara, M., Orihashi, Y., Yoneda, S., Shimizu, H., Kunimaru, T., Takahashi, K., Yanagi, T., Nakano, T., Fujimaki, H., Shinjo, R., Asahara, Y., Tanimizu, M., Dragusanu, C., 2000. JNdi-1: a neodymium isotopic reference in consistency with LaJolla neodymium. *Chem. Geol.* 168, 279–281.
- Taylor, R.N., Ishizuka, O., Michalik, A., Milton, J.A., Croudace, I.W., 2015. Evaluating the precision of Pb isotope measurement by mass spectrometry. *J. Anal. Atomic Spectrom.* 30, 198–213.
- Tchakerian, V.P., Lancaster, N., 2002. Late quaternary arid/humid cycles in the Mojave Desert and western Great Basin of North America. *Quat. Sci. Rev.* 21, 799–810.
- Tiedemann, R., Sarnthein, M., Shackleton, N.J., 1994. Astronomic timescale for the pliocene Atlantic $\delta^{18}O$ and dust flux records of Ocean Drilling Program site 659. *Paleoceanography* 9, 619–638.
- Toomey, M., 2010. KML to SHP conversion using kml_shapefile, 1.2.0.0. In: MATLAB Central File Exchange.
- Uno, I., Eguchi, K., Yumimoto, K., Liu, Z., Hara, Y., Sugimoto, N., Shimizu, A., Takemura, T., 2011. Large Asian dust layers continuously reached North America in April 2010. *Atmos. Chem. Phys.* 11, 7333–7341.
- Urban, F.E., Goldstein, H.L., Fulton, R., Reynolds, R.L., 2018. Unseen dust emission and global dust abundance: documenting dust emission from the Mojave Desert (USA) by daily remote camera imagery and wind-erosion measurements. *J. Geophys. Res.-Atmospheres* 123, 8735–8753.
- van der Does, M., Knippertz, P., Zschenderlein, P., Giles Harrison, R., Stuut, J.-B.W., 2018a. The mysterious long-range transport of giant mineral dust particles. *Sci. Adv.* 4.
- van der Does, M., Pourmand, A., Sharifi, A., Stuut, J.-B.W., 2018b. North African mineral dust across the tropical Atlantic Ocean: insights from dust particle size, radiogenic Sr-Nd-Hf isotopes and rare earth elements (REE). *Aeolian Res.* 33, 106–116.
- Vance, D., Thirlwall, M., 2002. An assessment of mass discrimination in MC-ICPMS using Nd isotopes. *Chem. Geol.* 185, 227–240.
- VanCuren, R.A., Cahill, T.A., 2002. Asian aerosols in north America: frequency and concentration of fine dust. *J. Geophys. Res. Atmosp.* 107, AAC 19–11–AAC 19–16.
- Walter, H.J., Hegner, E., Diekmann, B., Kuhn, G., Rutgers van der loeff, M.M., 2000. Provenance and transport of terrigenous sediment in the south Atlantic Ocean and their relations to glacial and interglacial cycles: Nd and Sr isotopic evidence. *Geochem. Cosmochim. Acta* 64, 3813–3827.
- Weigand, P., 1993. Geochemistry and Origin of Middle Miocene Volcanic Rocks from Santa Cruz and Anacapa Islands, Southern California Borderland. Santa Barbara Museum of Natural History, Santa Barbara, CA.
- Weigand, P.W., Savage, K.L., 1999. Summary of the Miocene igneous rocks of the Channel islands, southern California. In: Browne, D.R.H., Chaney, H., Mitchell (Eds.), *Fifth Californian Islands Symposium*, (Eds.).
- Wu, F., Ho, S.S.H., Sun, Q.L., Ip, S.H.S., 2011. Provenance of Chinese loess: evidence from stable lead isotope. *Terr. Atmos. Ocean Sci.* 22, 305–314.
- Xie, R.F.C., Marcantonio, F., 2012. Deglacial dust provenance changes in the Eastern Equatorial Pacific and implications for ITCZ movement. *Earth Planet Sci. Lett.* 317, 386–395.
- Zeng, F., Liang, M., Peng, S., Yu, D., Xiang, S., 2015. Sr-Nd-Pb isotopic compositions of the Neogene eolian deposits in the Xining Basin and implications for their dust sources. *J. Earth Sci.* 26, 669–676.
- Zhao, W., Sun, Y., Balsam, W., Zeng, L., Lu, H., Otgonbayar, K., Ji, J., 2015. Clay-sized Hf-Nd-Sr isotopic composition of Mongolian dust as a fingerprint for regional to hemispherical transport. *Geophys. Res. Lett.* 42, 5661–5669.